

Applicant : Alan D. Snow *et al.*
 Serial No. : 10/077,596
 Filed : February 15, 2002
 Page : 2 of 31


Attorney's Docket No.: 17170-006001

This listing of claims replaces all prior versions and listings of claims in the application:


Listing of Claims:

1-27. (Cancelled)

28. (Previously presented) A pharmaceutical composition comprising a therapeutically effective amount of a proanthocyanidin, selected from a group of proanthocyanidins characterized by Formula I or Formula II, and proanthocyanidins characterized by oligomeric combinations of Formula I and Formula II, and pharmaceutically acceptable salts of the foregoing proanthocyanidins:



Formula I



Formula II

where:

n is an integer of 2 to 20;

R_1 and R_2 are independently selected from hydrogen and hydroxy;

R_3 is selected from the group consisting of hydrogen, optionally substituted O-glycosyl, -C(O)-(optionally substituted aryl), and BC(O)-(optionally substituted heteroaryl);

R_4 is selected from the group consisting of hydrogen, catechin, epicatechin, epiafzelechin, and gallates of catechin and epicatechin;

the lines at the 2-, 3- and 4-position denote optional R and S configurations;

the lines at the 4- and 8-positions in Formula I and at the 4- and 6- positions in Formula II denote possible oligomer bonds between individual units, and

Applicant : Alan D. Snow *et al.*
Serial No. : 10/077,596
Filed : February 15, 2002
Page : 3 of 31

Attorney's Docket No.: 17170-006001

the substitutions at R₁, R₂, R₃, and R₄, and the configurations at the 2-, 3-, and 4-positions, and the oligomer bond configurations of 4-8 and 4-6 are independently selected for each individual unit and a pharmaceutically acceptable carrier, diluent, or excipient, the therapeutic amount of the proanthocyanidin selected for efficacy in treating amyloid, α -synuclein or NAC fibrillogenesis in a mammalian subject.

29. (Previously presented) The composition of claim 28, wherein the therapeutically effective amount of the proanthocyanidin comprises a dosage in the range of about 10 to 1,000 mg/kg of body weight of the subject.

30. (Previously presented) The composition of claim 29, wherein the therapeutically effective amount of the proanthocyanidin comprises a dosage in the range of about 10 to 100 mg/kg of body weight of the subject.

31. (Previously presented) The composition of claim 29, wherein the proanthocyanidin is selected from the group consisting of dimers and trimers of epicatechin, epiafzelechin and catechin, and the pharmaceutically acceptable salts thereof.

32. (Previously presented) The composition of claim 31, wherein the proanthocyanidin is the procyanidin dimer epicatechin-4 β →8-epicatechin.

33. (Previously presented) The composition of claim 31, wherein the proanthocyanidin is the procyanidin dimer catechin-4 α →8-epicatechin.

34. (Previously presented) The composition of claim 31, wherein the proanthocyanidin is the procyanidin dimer epiafzelechin-4 β →8-epicatechin.

35. (Previously presented) The composition of claim 31, wherein the proanthocyanidin is the procyanidin trimer epicatechin-4 β →8-epicatechin-4 β →8-epicatechin.

36. (Previously presented) The composition of claim 31 comprising a mixture of two or more of the proanthocyanidins selected from the group consisting of dimers and trimers of epicatechin, epiafzelechin and catechin, and the pharmaceutically acceptable salts thereof.

37. (Previously presented) The composition of claim 36 comprising a mixture of two or more of the procyanidins selected from the group consisting of the dimers and trimers of epicatechin, and the pharmaceutically acceptable salts thereof.

Applicant : Alan D. Snow *et al.*
 Serial No. : 10/077,596
 Filed : February 15, 2002
 Page : 4 of 31

Attorney's Docket No.: 17170-006001

38. (Previously presented) The composition of claim 36 comprising a mixture of two or more of the proanthocyanidins selected from the group consisting of epicatechin-4 β →8-epicatechin, catechin-4 α →8-epicatechin, epiafzelechin-4 β →8-epicatechin, and epicatechin-4 β →8-epicatechin-4 β →8-epicatechin.


39. (Previously presented) The composition of claim 31, wherein each proanthocyanidin selected is present in a percentage purity that significantly exceeds a proportion percentage of the proanthocyanidin presence in a plant, or extract from a plant.

40. (Previously presented) The composition of claim 39, wherein the proanthocyanidin selected is at least a 70% pure proanthocyanidin.


41. (Previously presented) The composition of claim 40, wherein the proanthocyanidin selected is in at least 70% pure isolated or synthetic form.

42-54. (Cancelled)

55. (Previously presented) A pharmaceutical composition comprising a therapeutically effective amount of a mixture of at least 70% pure proanthocyanidins, selected from a group of proanthocyanidins characterized by Formula I or Formula II, and proanthocyanidins characterized by oligomeric combinations of Formula I and Formula II, and pharmaceutically acceptable salts of the foregoing proanthocyanidins:



Formula I



Formula II

where:

n is an integer of 2 to 20;

R₁ and R₂ are independently selected from hydrogen and hydroxy;

" Applicant : Alan D. Snow *et al.*
Serial No. : 10/077,596
Filed : February 15, 2002
Page : 5 of 31

Attorney's Docket No.: 17170-006001

R₃ is selected from the group consisting of hydrogen, optionally substituted O-glycosyl, -C(O)-(optionally substituted aryl), and BC(O)-(optionally substituted heteroaryl);
R₄ is selected from the group consisting of hydrogen, catechin, epicatechin, epiafzelechin, and gallates of catechin and epicatechin;
the lines at the 2-, 3- and 4-position denote optional R and S configurations;
the lines at the 4- and 8-positions in Formula I and at the 4- and 6- positions in Formula II denote possible oligomer bonds between individual units, and
the substitutions at R₁, R₂, R₃, and R₄, and the configurations at the 2-, 3-, and 4-positions, and the oligomer bond configurations of 4-8 and 4-6 are independently selected for each individual unit.

56. (Previously presented) The composition of claim 55, wherein one or more of the proanthocyanidins are selected from the group consisting of the dimers and trimers of epicatechin, epiafzelechin, and catechin, and the pharmaceutically acceptable salts thereof.